

**Cdk1 / p34cdc2 Antibody - With BSA and Azide**  
**Mouse Monoclonal Antibody [Clone CDK1/873 ]**  
**Catalog # AH12825**

### Specification

#### Cdk1 / p34cdc2 Antibody - With BSA and Azide - Product Information

|                   |                             |
|-------------------|-----------------------------|
| Application       | IHC, IF, FC                 |
| Primary Accession | <a href="#">P06493</a>      |
| Other Accession   | <a href="#">983, 334562</a> |
| Reactivity        | Human, Monkey, Bovine       |
| Host              | Mouse                       |
| Clonality         | Monoclonal                  |
| Isotype           | Mouse / IgG2a, kappa        |
| Calculated MW     | 34kDa KDa                   |

#### Cdk1 / p34cdc2 Antibody - With BSA and Azide - Additional Information

##### Gene ID 983

##### Other Names

Cyclin-dependent kinase 1, CDK1, 2.7.11.22, 2.7.11.23, Cell division control protein 2 homolog, Cell division protein kinase 1, p34 protein kinase, CDK1, CDC2, CDC28A, CDKN1, P34CDC2

##### Application Note

<span class ="dilution\_IHC">IHC~~1:100~500</span><br /><span class ="dilution\_IF">IF~~1:50~200</span><br /><span class ="dilution\_FC">FC~~1:10~50</span>

##### Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

##### Precautions

Cdk1 / p34cdc2 Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

#### Cdk1 / p34cdc2 Antibody - With BSA and Azide - Protein Information

##### Name CDK1

##### Synonyms CDC2, CDC28A, CDKN1, P34CDC2

##### Function

Plays a key role in the control of the eukaryotic cell cycle by modulating the centrosome cycle as well as mitotic onset; promotes G2-M transition via association with multiple interphase cyclins (PubMed:<a href="http://www.uniprot.org/citations/16407259" target="\_blank">16407259</a>, PubMed:<a href="http://www.uniprot.org/citations/16933150" target="\_blank">16933150</a>, PubMed:<a href="http://www.uniprot.org/citations/17459720" target="\_blank">17459720</a>, PubMed:<a href="http://www.uniprot.org/citations/18356527" target="\_blank">18356527</a>, PubMed:<a href="http://www.uniprot.org/citations/19509060" target="\_blank">19509060</a>,

PubMed:<a href="http://www.uniprot.org/citations/19917720" target="\_blank">19917720</a>,  
PubMed:<a href="http://www.uniprot.org/citations/20171170" target="\_blank">20171170</a>,  
PubMed:<a href="http://www.uniprot.org/citations/20935635" target="\_blank">20935635</a>,  
PubMed:<a href="http://www.uniprot.org/citations/20937773" target="\_blank">20937773</a>,  
PubMed:<a href="http://www.uniprot.org/citations/21063390" target="\_blank">21063390</a>,  
PubMed:<a href="http://www.uniprot.org/citations/2188730" target="\_blank">2188730</a>,  
PubMed:<a href="http://www.uniprot.org/citations/23355470" target="\_blank">23355470</a>,  
PubMed:<a href="http://www.uniprot.org/citations/2344612" target="\_blank">2344612</a>,  
PubMed:<a href="http://www.uniprot.org/citations/23601106" target="\_blank">23601106</a>,  
PubMed:<a href="http://www.uniprot.org/citations/23602554" target="\_blank">23602554</a>,  
PubMed:<a href="http://www.uniprot.org/citations/25556658" target="\_blank">25556658</a>,  
PubMed:<a href="http://www.uniprot.org/citations/26829474" target="\_blank">26829474</a>,  
PubMed:<a href="http://www.uniprot.org/citations/27814491" target="\_blank">27814491</a>,  
PubMed:<a href="http://www.uniprot.org/citations/30139873" target="\_blank">30139873</a>,  
PubMed:<a href="http://www.uniprot.org/citations/30704899" target="\_blank">30704899</a>).  
Phosphorylates PARVA/actopaxin, APC, AMPH, APC, BARD1, Bcl-xL/BCL2L1, BRCA2, CALD1, CASP8,  
CDC7, CDC20, CDC25A, CDC25C, CC2D1A, CENPA, CSNK2 proteins/CKII, FZR1/CDH1, CDK7,  
CEBPB, CHAMP1, DMD/dystrophin, EEF1 proteins/EF-1, EZH2, KIF11/EG5, EGFR, FANCG, FOS, GFAP,  
GOLGA2/GM130, GRASP1, UBE2A/hHR6A, HIST1H1 proteins/histone H1, HMGA1, HIVEP3/KRC,  
KAT5, LMNA, LMNB, LBR, MKI67, LATS1, MAP1B, MAP4, MARCKS, MCM2, MCM4, MKLP1, MLST8,  
MYB, NEFH, NFIC, NPC/nuclear pore complex, PITPNM1/NIR2, NPM1, NCL, NUCKS1, NPM1/numatrin,  
ORC1, PRKAR2A, EEF1E1/p18, EIF3F/p47, p53/TP53, NONO/p54NRB, PAPOLA, PLEC/plectin, RB1,  
TPPP, UL40/R2, RAB4A, RAP1GAP, RBBP8/CtIP, RCC1, RPS6KB1/S6K1, KHDRBS1/SAM68, ESPL1, SKI,  
BIRC5/survivin, STIP1, TEX14, beta-tubulins, MAPT/TAU, NEDD1, VIM/vimentin, TK1, FOXO1,  
RUNX1/AML1, SAMHD1, SIRT2, CGAS and RUNX2 (PubMed:<a  
href="http://www.uniprot.org/citations/16407259" target="\_blank">16407259</a>, PubMed:<a  
href="http://www.uniprot.org/citations/16933150" target="\_blank">16933150</a>, PubMed:<a  
href="http://www.uniprot.org/citations/17459720" target="\_blank">17459720</a>, PubMed:<a  
href="http://www.uniprot.org/citations/18356527" target="\_blank">18356527</a>, PubMed:<a  
href="http://www.uniprot.org/citations/19202191" target="\_blank">19202191</a>, PubMed:<a  
href="http://www.uniprot.org/citations/19509060" target="\_blank">19509060</a>, PubMed:<a  
href="http://www.uniprot.org/citations/19917720" target="\_blank">19917720</a>, PubMed:<a  
href="http://www.uniprot.org/citations/20171170" target="\_blank">20171170</a>, PubMed:<a  
href="http://www.uniprot.org/citations/20935635" target="\_blank">20935635</a>, PubMed:<a  
href="http://www.uniprot.org/citations/20937773" target="\_blank">20937773</a>, PubMed:<a  
href="http://www.uniprot.org/citations/21063390" target="\_blank">21063390</a>, PubMed:<a  
href="http://www.uniprot.org/citations/2188730" target="\_blank">2188730</a>, PubMed:<a  
href="http://www.uniprot.org/citations/23355470" target="\_blank">23355470</a>, PubMed:<a  
href="http://www.uniprot.org/citations/2344612" target="\_blank">2344612</a>, PubMed:<a  
href="http://www.uniprot.org/citations/23601106" target="\_blank">23601106</a>, PubMed:<a  
href="http://www.uniprot.org/citations/23602554" target="\_blank">23602554</a>, PubMed:<a  
href="http://www.uniprot.org/citations/25012651" target="\_blank">25012651</a>, PubMed:<a  
href="http://www.uniprot.org/citations/25556658" target="\_blank">25556658</a>, PubMed:<a  
href="http://www.uniprot.org/citations/26829474" target="\_blank">26829474</a>, PubMed:<a  
href="http://www.uniprot.org/citations/27814491" target="\_blank">27814491</a>, PubMed:<a  
href="http://www.uniprot.org/citations/30704899" target="\_blank">30704899</a>, PubMed:<a  
href="http://www.uniprot.org/citations/32351706" target="\_blank">32351706</a>, PubMed:<a  
href="http://www.uniprot.org/citations/34741373" target="\_blank">34741373</a>).  
CDK1/CDC2-cyclin-B controls pronuclear union in interphase fertilized eggs (PubMed:<a  
href="http://www.uniprot.org/citations/18480403" target="\_blank">18480403</a>, PubMed:<a  
href="http://www.uniprot.org/citations/20360007" target="\_blank">20360007</a>). Essential for  
early stages of embryonic development (PubMed:<a  
href="http://www.uniprot.org/citations/18480403" target="\_blank">18480403</a>, PubMed:<a  
href="http://www.uniprot.org/citations/20360007" target="\_blank">20360007</a>). During G2  
and early mitosis, CDC25A/B/C-mediated dephosphorylation activates CDK1/cyclin complexes  
which phosphorylate several substrates that trigger at least centrosome separation, Golgi  
dynamics, nuclear envelope breakdown and chromosome condensation (PubMed:<a

href="http://www.uniprot.org/citations/18480403" target="\_blank">>18480403</a>, PubMed:<a href="http://www.uniprot.org/citations/20360007" target="\_blank">>20360007</a>, PubMed:<a href="http://www.uniprot.org/citations/2188730" target="\_blank">>2188730</a>, PubMed:<a href="http://www.uniprot.org/citations/2344612" target="\_blank">>2344612</a>, PubMed:<a href="http://www.uniprot.org/citations/30139873" target="\_blank">>30139873</a>). Once chromosomes are condensed and aligned at the metaphase plate, CDK1 activity is switched off by WEE1- and PKMYT1-mediated phosphorylation to allow sister chromatid separation, chromosome decondensation, reformation of the nuclear envelope and cytokinesis (PubMed:<a href="http://www.uniprot.org/citations/18480403" target="\_blank">>18480403</a>, PubMed:<a href="http://www.uniprot.org/citations/20360007" target="\_blank">>20360007</a>).

Phosphorylates KRT5 during prometaphase and metaphase (By similarity). Inactivated by PKR/EIF2AK2- and WEE1-mediated phosphorylation upon DNA damage to stop cell cycle and genome replication at the G2 checkpoint thus facilitating DNA repair (PubMed:<a href="http://www.uniprot.org/citations/20360007" target="\_blank">>20360007</a>). Reactivated after successful DNA repair through WIP1-dependent signaling leading to CDC25A/B/C-mediated dephosphorylation and restoring cell cycle progression (PubMed:<a href="http://www.uniprot.org/citations/20395957" target="\_blank">>20395957</a>). Catalyzes lamin (LMNA, LMNB1 and LMNB2) phosphorylation at the onset of mitosis, promoting nuclear envelope breakdown (PubMed:<a href="http://www.uniprot.org/citations/2188730" target="\_blank">>2188730</a>, PubMed:<a href="http://www.uniprot.org/citations/2344612" target="\_blank">>2344612</a>, PubMed:<a href="http://www.uniprot.org/citations/37788673" target="\_blank">>37788673</a>). In proliferating cells, CDK1-mediated FOXO1 phosphorylation at the G2-M phase represses FOXO1 interaction with 14-3-3 proteins and thereby promotes FOXO1 nuclear accumulation and transcription factor activity, leading to cell death of postmitotic neurons (PubMed:<a href="http://www.uniprot.org/citations/18356527" target="\_blank">>18356527</a>). The phosphorylation of beta-tubulins regulates microtubule dynamics during mitosis (PubMed:<a href="http://www.uniprot.org/citations/16371510" target="\_blank">>16371510</a>). NEDD1 phosphorylation promotes PLK1-mediated NEDD1 phosphorylation and subsequent targeting of the gamma-tubulin ring complex (gTuRC) to the centrosome, an important step for spindle formation (PubMed:<a href="http://www.uniprot.org/citations/19509060" target="\_blank">>19509060</a>). In addition, CC2D1A phosphorylation regulates CC2D1A spindle pole localization and association with SCC1/RAD21 and centriole cohesion during mitosis (PubMed:<a href="http://www.uniprot.org/citations/20171170" target="\_blank">>20171170</a>). The phosphorylation of Bcl-xL/BCL2L1 after prolonged G2 arrest upon DNA damage triggers apoptosis (PubMed:<a href="http://www.uniprot.org/citations/19917720" target="\_blank">>19917720</a>). In contrast, CASP8 phosphorylation during mitosis prevents its activation by proteolysis and subsequent apoptosis (PubMed:<a href="http://www.uniprot.org/citations/20937773" target="\_blank">>20937773</a>). This phosphorylation occurs in cancer cell lines, as well as in primary breast tissues and lymphocytes (PubMed:<a href="http://www.uniprot.org/citations/20937773" target="\_blank">>20937773</a>). EZH2 phosphorylation promotes H3K27me3 maintenance and epigenetic gene silencing (PubMed:<a href="http://www.uniprot.org/citations/20935635" target="\_blank">>20935635</a>). CALD1 phosphorylation promotes Schwann cell migration during peripheral nerve regeneration (By similarity). CDK1-cyclin-B complex phosphorylates NCKAP5L and mediates its dissociation from centrosomes during mitosis (PubMed:<a href="http://www.uniprot.org/citations/26549230" target="\_blank">>26549230</a>). Regulates the amplitude of the cyclic expression of the core clock gene BMAL1 by phosphorylating its transcriptional repressor NR1D1, and this phosphorylation is necessary for SCF(FBXW7)-mediated ubiquitination and proteasomal degradation of NR1D1 (PubMed:<a href="http://www.uniprot.org/citations/27238018" target="\_blank">>27238018</a>). Phosphorylates EML3 at 'Thr-881' which is essential for its interaction with HAUS augmin-like complex and TUBG1 (PubMed:<a href="http://www.uniprot.org/citations/30723163" target="\_blank">>30723163</a>). Phosphorylates CGAS during mitosis, leading to its inhibition, thereby preventing CGAS activation by self DNA during mitosis (PubMed:<a href="http://www.uniprot.org/citations/32351706" target="\_blank">>32351706</a>). Phosphorylates SKA3 on multiple sites during mitosis which promotes SKA3 binding to the NDC80 complex and anchoring of the SKA complex to kinetochores, to enable stable attachment of mitotic spindle microtubules to kinetochores (PubMed:<a href="http://www.uniprot.org/citations/30723163" target="\_blank">>30723163</a>).

[28479321](http://www.uniprot.org/citations/28479321), PubMed:[31804178](http://www.uniprot.org/citations/31804178), PubMed:[32491969](http://www.uniprot.org/citations/32491969)).

#### Cellular Location

Nucleus {ECO:0000250|UniProtKB:P11440}. Cytoplasm {ECO:0000250|UniProtKB:P11440}. Mitochondrion. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Note=Cytoplasmic during the interphase Colocalizes with SIRT2 on centrosome during prophase and on spindle fibers during metaphase of the mitotic cell cycle. Reversibly translocated from cytoplasm to nucleus when phosphorylated before G2-M transition when associated with cyclin-B1. Accumulates in mitochondria in G2-arrested cells upon DNA-damage

#### Tissue Location


[Isoform 2]: Found in breast cancer tissues.

#### Cdk1 / p34cdc2 Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Cdk1 / p34cdc2 Antibody - With BSA and Azide - Images



Formalin-fixed, paraffin-embedded human Tonsil stained with Cdk1 Monoclonal Antibody (CDK1/873).

#### Cdk1 / p34cdc2 Antibody - With BSA and Azide - Background

Recognizes a 34kDa protein, which is identified as cyclin dependent kinase 1 (cdk1) or p34cdc2 protein kinase. cdk1 / p34cdc2 plays a crucial role during cell division and is most active during mitosis. It is predominantly localized in the nucleus. It is a serine/threonine kinase, which is activated by cyclin, presumably by de-phosphorylation of tyrosine residues. Activated cdk1 / p34cdc2 performs specific functions during mitosis, including nuclear envelope breakdown and

chromosome condensation.

**Cdk1 / p34cdc2 Antibody - With BSA and Azide - References**

Lukas J et. al. European Journal of Biochemistry, 1992, 207(1):169-76