

**Lewis A (Blood Group Antigen) Antibody - With BSA and Azide**  
**Mouse Monoclonal Antibody [Clone SPM522 ]**  
**Catalog # AH12954**

## Specification

---

### Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - Product Information

Application	,2,3,4,
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	Multiple KDa

### Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - Additional Information

#### Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

#### Precautions

Lewis A (Blood Group Antigen) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.


### Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - Protein Information


### Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - Images





Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with Lewis A Monoclonal Antibody (SPM522).

#### **Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - Background**

Recognizes a carbohydrate determinant of Gal 1-3(Fuc 1-4) GlcNAc which is blood group antigen Lewis A. It is present primarily on epithelial cells such as colon and kidneys. In the tumors and dedifferentiated tissues, decrease of Lewis A antigen was observed. Lewis A (type 1 chain) is expressed in colonic epithelial cells and may be useful for detection of gastrointestinal tumors, pancreatic cancer, and colorectal tumors. Blood group related antigens represent a group of carbohydrate determinants carried on both glycolipids and glycoproteins. They are usually mucin-type, and are detected on erythrocytes, certain epithelial cells, and in secretions of certain individuals. Sixteen genetically and biosynthetically distinct but inter-related specificities belong to this group of antigens, including A, B, H, Lewis A, Lewis B, Lewis X, Lewis Y, and precursor type 1 chain antigens.

#### **Lewis A (Blood Group Antigen) Antibody - With BSA and Azide - References**

Blood transfusion and immunohaematology, Ph. Rouger, D Anstee and Ch. Salmon (Eds). Arnette, France 30 (5): 353-720 (1987). | Cancer Epidemiology, Biomarkers Prevention 1, 199-205 (1992). | Gastroenterology, 102 (2), 424-430 (1992). |