

Mitochondria (Marker for Human Cells) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone AE-1]
Catalog # AH13042

Specification

Mitochondria (Marker for Human Cells) Antibody - With BSA and Azide - Product Information

Application	,1,2,3,10,
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	60kDa KDa

Mitochondria (Marker for Human Cells) Antibody - With BSA and Azide - Additional Information

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

Mitochondria (Marker for Human Cells) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.


Mitochondria (Marker for Human Cells) Antibody - With BSA and Azide - Protein Information


Mitochondria (Marker for Human Cells) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mitochondria (Marker for Human Cells) Antibody - With BSA and Azide - Images





Formalin-fixed, paraffin-embedded human Tonsil stained with Mitochondria Monoclonal Antibody (AE-1).



Formalin-fixed, paraffin-embedded human Pancreas stained with Mitochondria Monoclonal Antibody (AE-1).

Mitochondria (Marker for Human Cells) Antibody - With BSA and Azide - Background

This MAb recognizes a 60kDa antigen associated with the mitochondria in human cells. It is a part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. It recognizes an antigen associated with the mitochondria in human cells only. It can be used to stain the mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells and may be used to stain mitochondria of cells in fixed or frozen tissue sections. It can also be used with paraformaldehyde fixed frozen tissue or cell preparations. This MAb is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells, including neurons and embryonic stem cells.