

Anti-LH, alpha (Luteinizing Hormone, alpha) Antibody Mouse Monoclonal Antibody Catalog # AH13107

Specification

Anti-LH, alpha (Luteinizing Hormone, alpha) Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW IHC-P <u>P01215</u> <u>119689</u> Human Mouse Monoclonal Mouse / IgG1, kappa 13075

Anti-LH, alpha (Luteinizing Hormone, alpha) Antibody - Additional Information

Gene ID 1081

Other Names

CG-alpha; CGA; Chorionic Gonadotrophin Alpha; Follicle Stimulating Hormone Alpha; Follitropin Alpha; FSH-alpha; FSHA; GPH Alpha; GPHA1; LHA; LH-alpha; Luteinizing Hormone Alpha; Lutropin Alpha; Thyroid Stimulating Hormone Alpha; Thyrotropin Alpha; TSHA

Application Note IHC-P~~N/A

Format

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage

Store at 2 to 8°C.Antibody is stable for 24 months.

Precautions

Anti-LH, alpha (Luteinizing Hormone, alpha) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-LH, alpha (Luteinizing Hormone, alpha) Antibody - Protein Information

Name CGA

Function

Shared alpha chain of the active heterodimeric glycoprotein hormones thyrotropin/thyroid stimulating hormone/TSH, lutropin/luteinizing hormone/LH, follitropin/follicle stimulating hormone/FSH and choriogonadotropin/CG. These hormones bind specific receptors on target cells that in turn activate downstream signaling pathways.



Cellular Location Secreted

Anti-LH, alpha (Luteinizing Hormone, alpha) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-LH, alpha (Luteinizing Hormone, alpha) Antibody - Images



Formalin-fixed, paraffin-embedded human Pituitary stained with LH alpha Monoclonal Antibody (LHa/756).

Anti-LH, alpha (Luteinizing Hormone, alpha) Antibody - Background

This MAb reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunits vary. LH has a beta subunit of 121 amino acids (LHB) that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunit contains the same amino acids in sequence as the beta subunit of hCG and both stimulate the same receptor; however, the hCG beta subunit contains an additional 24 amino acids and the hormones differ in the composition of their sugar moieties. LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other pituitary gonadotropin follicle-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary tumors and the study of pituitary disease.