

**Anti-Desmoglein-2 (DSG2) Antibody**  
**Mouse Monoclonal Antibody**  
**Catalog # AH13172****Specification**

---

**Anti-Desmoglein-2 (DSG2) Antibody - Product Information**

Application	WB, IF, FC
Primary Accession	<a href="#">Q14126</a>
Other Accession	<a href="#">412597</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	122294

**Anti-Desmoglein-2 (DSG2) Antibody - Additional Information****Gene ID** 1829**Other Names**ARVC10; ARVD10; Cadherin family member 5; CDHF5; CMD1BB; Desmoglein-2; DSG2; HDGC;  
Human Desmoglein Colon**Application Note**

WB~~1:1000  
IF~~1:50~200  
FC~~1:10~50

**Format**

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA &amp; 0.05% azide. Also available WITHOUT BSA &amp; azide at 1.0mg/ml.

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

Anti-Desmoglein-2 (DSG2) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-Desmoglein-2 (DSG2) Antibody - Protein Information****Name** DSG2**Synonyms** CDHF5**Function**A component of desmosome cell-cell junctions which are required for positive regulation of cellular adhesion (PubMed: <http://www.uniprot.org/citations/38395410> target="\_blank">38395410</a>). Involved in the interaction of plaque proteins and intermediate

filaments mediating cell-cell adhesion. Required for proliferation and viability of embryonic stem cells in the blastocyst, thereby crucial for progression of post-implantation embryonic development (By similarity). Maintains pluripotency by regulating epithelial to mesenchymal transition/mesenchymal to epithelial transition (EMT/MET) via interacting with and sequestering CTNNB1 to sites of cell-cell contact, thereby reducing translocation of CTNNB1 to the nucleus and subsequent transcription of CTNNB1/TCF-target genes (PubMed:<a href="http://www.uniprot.org/citations/29910125" target="\_blank">29910125</a>). Promotes pluripotency and the multi-lineage differentiation potential of hematopoietic stem cells (PubMed:<a href="http://www.uniprot.org/citations/27338829" target="\_blank">27338829</a>). Plays a role in endothelial cell sprouting and elongation via mediating the junctional-association of cortical actin fibers and CDH5 (PubMed:<a href="http://www.uniprot.org/citations/27338829" target="\_blank">27338829</a>). Plays a role in limiting inflammatory infiltration and the apoptotic response to injury in kidney tubular epithelial cells, potentially via its role in maintaining cell-cell adhesion and the epithelial barrier (PubMed:<a href="http://www.uniprot.org/citations/38395410" target="\_blank">38395410</a>).

### **Cellular Location**

Cell membrane; Single-pass type I membrane protein. Cell junction, desmosome. Cytoplasm

### **Tissue Location**

Expressed in undifferentiated pluripotent stem cells, expression decreases during differentiation (at protein level) (PubMed:29910125). Expressed in hematopoietic stem cells and circulating endothelial progenitor cells, expression decreases upon increasing cell lineage commitment (at protein level) (PubMed:27338829). Expressed on common myeloid progenitors, pro- myelocytes, pro-erythrocytes and B-cell lineage progenitors (at protein level). Expression in mature cell types in the bone marrow and mature leukocyte populations is absent (PubMed:27338829). Expressed by foreskin fibroblasts, expression peaks during the early stage of differentiation reprogramming (at protein level) (PubMed:29910125) Expressed by endothelial cells in both arterioles and venules in the cervix (at protein level) (PubMed:27338829). Expressed in kidney tubular epithelial cells (PubMed:38395410)

## **Anti-Desmoglein-2 (DSG2) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Anti-Desmoglein-2 (DSG2) Antibody - Images**

## **Anti-Desmoglein-2 (DSG2) Antibody - Background**

Recognizes a protein of 165kDa, identified as Desmoglein-2 (DSG2). This monoclonal antibody recognizes the extracellular domain of human desmoglein-2. Desmoglein-2 is a member of the desmosomal cadherin family. Desmosomes are intercellular adhering junctions that represent cell surface attachment sites for intermediate filament. Desmocollins and desmogleins are the main desmosomal transmembrane proteins. Desmogleins consist of Dsg1, Dsg2, Dsg3, and Dsg4 isoforms. Within the desmosome, the extracellular domain of desmoglein is essential for calcium dependent heterophilic binding to the desmocollins, whereas the intracellular domain is essential for binding to the desmosomal plaque protein, plakoglobin. Human Desmoglein-2 is a type I

transmembrane glycoprotein of 1117 amino acid (aa) residues with a 23 aa signal peptide and a 25 aa propeptide. It differs from other classic cadherins by having four instead of five cadherin repeat domains in its extracellular region, and a much larger cytoplasmic region containing five desmoglein repeat domains which share homology with the cadherin repeats. Instead of having the HAV adhesion motif found in type I cadherins, desmogleins have R/YAL as the adhesion motif on its amino-terminal cadherin repeat. The cytoplasmic tails of desmogleins interact with desmoplakins, plakoglobin and plakophilins. In turn, these proteins link the desmogleins with the intermediate filaments. Desmoglein-2 has been shown to be important in establishing cell-cell adhesion and function in epithelial cells. Desmoglein2 was originally identified in colon carcinoma and colon, and was named HDGC (human desmoglein colon).