

**Anti-AMACR / p504S (Prostate Cancer Marker) Antibody**  
**Mouse Monoclonal Antibody**  
**Catalog # AH13243****Specification****Anti-AMACR / p504S (Prostate Cancer Marker) Antibody - Product Information**

Application	IHC-P, IF
Primary Accession	<a href="#">O9UHK6</a>
Other Accession	<a href="#">508343</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG
Calculated MW	42387

**Anti-AMACR / p504S (Prostate Cancer Marker) Antibody - Additional Information****Gene ID** 23600**Other Names**

Alpha-methylacyl-CoA Racemase, CBAS4, Da1-8, Macr1, RACE, RM

**Application Note**

IHC-P~N/A  
IF~1:50~200

**Format**

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA &amp; 0.05% azide. Also available WITHOUT BSA &amp; azide at 1.0mg/ml.

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

Anti-AMACR / p504S (Prostate Cancer Marker) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-AMACR / p504S (Prostate Cancer Marker) Antibody - Protein Information****Name** AMACR**Function**

Catalyzes the interconversion of (R)- and (S)-stereoisomers of alpha-methyl-branched-chain fatty acyl-CoA esters (PubMed: [10655068](http://www.uniprot.org/citations/10655068), PubMed: [11060359](http://www.uniprot.org/citations/11060359), PubMed: [7649182](http://www.uniprot.org/citations/7649182)). Acts only on coenzyme A thioesters, not on free fatty acids, and accepts as substrates a wide range of alpha-methylacyl-CoAs, including pristanoyl-CoA,

trihydroxycoprostanoyl-CoA (an intermediate in bile acid synthesis), and arylpropionic acids like the anti-inflammatory drug ibuprofen (2- (4-isobutylphenyl)propionic acid) but neither 3-methyl-branched nor linear-chain acyl-CoAs (PubMed:<a href="http://www.uniprot.org/citations/10655068" target="\_blank">10655068</a>, PubMed:<a href="http://www.uniprot.org/citations/11060359" target="\_blank">11060359</a>, PubMed:<a href="http://www.uniprot.org/citations/7649182" target="\_blank">7649182</a>).

#### Cellular Location


Peroxisome. Mitochondrion

### Anti-AMACR / p504S (Prostate Cancer Marker) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-AMACR / p504S (Prostate Cancer Marker) Antibody - Images



Formalin-fixed, paraffin-embedded human Prostate Carcinoma Stained with AMACR / p504S Monoclonal Antibody (AMACR/1723)

### Anti-AMACR / p504S (Prostate Cancer Marker) Antibody - Background

This antibody recognizes a protein of 54kDa, which is identified as AMACR, also known as p504S. It is an enzyme that is involved in bile acid biosynthesis and  $\beta$ -oxidation of branched-chain fatty acids. AMACR is essential in lipid metabolism. It is expressed in cells of premalignant high-grade prostatic intraepithelial neoplasia (HGPIN) and prostate adenocarcinoma. The majority of the carcinoma cells show a distinct granular cytoplasmic staining reaction. AMACR is present at low or undetectable levels in glandular epithelial cells of normal prostate and benign prostatic hyperplasia. A spotty granular cytoplasmic staining is seen in a few cells of the benign glands. AMACR is expressed in normal liver (hepatocytes), kidney (tubular epithelial cells) and gall bladder (epithelial cells). Expression has also been found in lung (bronchial epithelial cells) and colon (colonic surface epithelium). AMACR expression can also be found in hepatocellular carcinoma and kidney carcinoma. Past studies have also shown that AMACR is expressed in various colon carcinomas

(well, moderately and poorly differentiated) and over expressed in prostate carcinoma.