

Anti-Cytokeratin 7 Antibody

Recombinant Mouse Monoclonal Antibody Catalog # AH13345

Specification

Anti-Cytokeratin 7 Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW IHC-P, IF, FC <u>P08729</u> <u>411501</u> Human Mouse Monoclonal Mouse / IgG1 51386

Anti-Cytokeratin 7 Antibody - Additional Information

Gene ID 3855

Other Names CK-7, K2C7, Keratin 55K Type II Cytoskeletal, Keratin Simple Epithelial Type 1 K7, Keratin Type II Cytoskeletal 7, Krt2-7, KRT7, Sarcolectin, SCL, Type II Mesothelial Keratin K7, Type-II Keratin Kb7

Application Note IHC-P~~N/A<br \>IF~~1:50~200<br \>FC~~1:10~50

Format

200ug/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage

Store at 2 to 8°C.Antibody is stable for 24 months.

Precautions

Anti-Cytokeratin 7 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Cytokeratin 7 Antibody - Protein Information

Name KRT7

Synonyms SCL

Function

Blocks interferon-dependent interphase and stimulates DNA synthesis in cells. Involved in the translational regulation of the human papillomavirus type 16 E7 mRNA (HPV16 E7).



Cellular Location Cytoplasm.

Tissue Location

Expressed in cultured epidermal, bronchial and mesothelial cells but absent in colon, ectocervix and liver. Observed throughout the glandular cells in the junction between stomach and esophagus but is absent in the esophagus.

Anti-Cytokeratin 7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Cytokeratin 7 Antibody - Images



Formalin-fixed, paraffin-embedded human Endometrial Carcinoma stained with Cytokeratin-7 Recombinant Mouse Monoclonal Antibody (rOV-TL12/30).

Anti-Cytokeratin 7 Antibody - Background

It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This MAb is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This MAb is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-).