

Anti-ACTH (C-Terminus) Antibody

Mouse Monoclonal Antibody Catalog # AH13444

Specification

Anti-ACTH (C-Terminus) Antibody - Product Information

Application IF, FC, E
Primary Accession P01189
Other Accession 1897
Reactivity Human
Host Mouse
Clonality Monoclonal
Isotype Mouse / IgG1

Calculated MW 29424

Anti-ACTH (C-Terminus) Antibody - Additional Information

Gene ID 5443

Other Names

Adrenocorticotropin; alpha or beta or gamma Melanocyte Stimulating Hormone (MSH) or Melanotropin; beta-Endorphin; beta or gamma Lipotropin (LPH); CLIP; Met Enkephalin; POC; POMC

Application Note

```
<span class ="dilution_IF">IF~~1:50~200</span><br \> < span class
="dilution FC">FC~~1:10~50</span> < br \> < span class = "dilution E">E~~N/A</span>
```

Format

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

Anti-ACTH (C-Terminus) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-ACTH (C-Terminus) Antibody - Protein Information

Name POMC

Function

[Corticotropin]: Stimulates the adrenal glands to release cortisol. [Melanocyte-stimulating hormone beta]: Increases the pigmentation of skin by increasing melanin production in melanocytes. [Met-enkephalin]: Endogenous opiate.

Cellular Location



Tel: 858.875.1900 Fax: 858.875.1999

Secreted {ECO:0000250|UniProtKB:P01193}. Note=Melanocyte-stimulating hormone alpha and beta-endorphin are stored in separate granules in hypothalamic POMC neurons, suggesting that secretion may be under the control of different regulatory mechanisms {ECO:0000250|UniProtKB:P01193}

Tissue Location

ACTH and MSH are produced by the pituitary gland.

Anti-ACTH (C-Terminus) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-ACTH (C-Terminus) Antibody - Images

Anti-ACTH (C-Terminus) Antibody - Background

ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by the anterior pituitary. This MAb is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro-opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslational processing by convertases. POMC is cleaved into ten hormone chains named NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gamma-MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotropin-gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells of immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituitary tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs). It also may react with other tumors (e.g. some small cell carcinomas of the lung) causing paraneoplastic syndromes by secreting ACTH.