

TLE1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10052**Specification**

TLE1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q04724
Other Accession	Q04724 , NP_005068 , NM_005077
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Rabbit, Pig, Chicken, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	83 kDa KDa

TLE1 antibody - N-terminal region - Additional Information**Gene ID** 7088**Alias Symbol** **ESG, ESG1, GRG1****Other Names**

Transducin-like enhancer protein 1, E(Sp1) homolog, Enhancer of split groucho-like protein 1, ESG1, TLE1

Target/Specificity

TLE1 is a transcriptional corepressor that binds to a number of transcription factors. TLE1 inhibits NF-kappa-B-regulated gene expression and the transcriptional activation mediated by FOXA2, and by CTNNB1 and TCF family members in Wnt signaling. The effects of full-length TLE family members may be modulated by association with dominant-negative AES. TLE1 has an unusual function as coactivator for ESRRG.

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-TLE1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

TLE1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TLE1 antibody - N-terminal region - Protein Information**Name** TLE1

Function

Transcriptional corepressor that binds to a number of transcription factors. Inhibits NF-kappa-B-regulated gene expression. Inhibits the transcriptional activation mediated by FOXA2, and by CTNNB1 and TCF family members in Wnt signaling. Enhances FOXG1/BF- 1- and HES1-mediated transcriptional repression (By similarity). The effects of full-length TLE family members may be modulated by association with dominant-negative AES. Unusual function as coactivator for ESRRG.

Cellular Location

Nucleus. Note=Nuclear and chromatin-associated, depending on isoforms and phosphorylation status. Hyperphosphorylation decreases the affinity for nuclear components


Tissue Location

In all tissues examined, mostly in brain, liver and muscle

TLE1 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TLE1 antibody - N-terminal region - Images

TLE1 antibody - N-terminal region (AI10052) in Human PANC1 cells using Western Blot
WB Suggested Anti-TLE1 Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:312500
Positive Control: PANC1 cell lysate

TLE1 antibody - N-terminal region - Background

This is a rabbit polyclonal antibody against TLE1. It was validated on Western Blot using a cell lysate as a positive control. Abgent strives to provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).