

DLX1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10073**Specification**

DLX1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P56177
Other Accession	P56177 , NP_835221 , NM_178120
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Chicken, Dog, Guinea Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27 kDa KDa

DLX1 antibody - C-terminal region - Additional Information**Gene ID** 1745**Other Names**

Homeobox protein DLX-1, DLX1

Target/Specificity

DLX1 is a member of a homeobox transcription factor family. It is localized to the nucleus where it may function as a transcriptional regulator of signals from multiple TGF- β superfamily members. DLX1 may play a role in the control of craniofacial patterning and the differentiation and survival of inhibitory neurons in the forebrain. This gene encodes a member of a homeobox transcription factor gene family similar to the Drosophila distal-less gene. The encoded protein is localized to the nucleus where it may function as a transcriptional regulator of signals from multiple TGF- β superfamily members. The encoded protein may play a role in the control of craniofacial patterning and the differentiation and survival of inhibitory neurons in the forebrain. This gene is located in a tail-to-tail configuration with another member of the family on the long arm of chromosome 2. Alternatively spliced transcript variants encoding different isoforms have been described.

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-DLX1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

DLX1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DLX1 antibody - C-terminal region - Protein Information

Name DLX1

Function

Plays a role as a transcriptional activator or repressor (PubMed:14671321). Inhibits several cytokine signaling pathways, such as TGF β 1, activin-A/INHBA and BMP4 by interfering with the transcriptional stimulatory activity of transcription factors, such as MSX2, FAST2, SMAD2 and SMAD3 during hematopoietic cell differentiation (PubMed:14671321). Plays a role in terminal differentiation of interneurons, such as amacrine and bipolar cells in the developing retina (By similarity). Likely to play a regulatory role in the development of the ventral forebrain (By similarity). May play a role in craniofacial patterning and morphogenesis and may be involved in the early development of diencephalic subdivisions (By similarity).

Cellular Location

Nucleus.

Tissue Location


Expressed in hematopoietic cell lines.

DLX1 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DLX1 antibody - C-terminal region - Images



DLX1 antibody - C-terminal region (AI10073) in Human Placenta cells using Western Blot
WB Suggested Anti-DLX1 Antibody Titration: 0.2-1 μ g/ml
ELISA Titer: 1:1562500
Positive Control: Human Placenta

DLX1 antibody - C-terminal region - Background

This is a rabbit polyclonal antibody against DLX1. It was validated on Western Blot using a cell lysate as a positive control. Abgent strives to provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).