

BRD3 Antibody - middle region

Rabbit Polyclonal Antibody Catalog # Al10087

Specification

BRD3 Antibody - middle region - Product Information

Application WB
Primary Accession O15059
Other Accession O15059-2
Reactivity Human

Predicted Human, Mouse, Rat, Rabbit, Zebrafish, Pig,

Dog, Guinea Pig, Horse, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 61 kDa KDa

BRD3 Antibody - middle region - Additional Information

Gene ID 8019

Other Names

Bromodomain-containing protein 3, RING3-like protein, BRD3, KIAA0043, RING3L

Target/Specificity

This gene was identified based on its homology to the gene encoding the RING3 protein, a serine/threonine kinase. The gene localizes to 9q34, a region which contains several major histocompatibility complex (MHC) genes. The function of the encoded protein is not known.

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul, I of distilled water. Final Anti-BRD3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions


BRD3 Antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

BRD3 Antibody - middle region - Protein Information

Name BRD3 {ECO:0000303|PubMed:18406326, ECO:0000312|HGNC:HGNC:1104}

Function

Chromatin reader that recognizes and binds acetylated histones, thereby controlling gene expression and remodeling chromatin structures (PubMed:18406326, PubMed:22464331, PubMed:<a





href="http://www.uniprot.org/citations/27105114" target="_blank">27105114, PubMed:32895492). Recruits transcription factors and coactivators to target gene sites, and activates RNA polymerase II machinery for transcriptional elongation (PubMed:29567837, PubMed:32895492). In vitro, binds acetylated lysine residues on the N-terminus of histone H2A, H2B, H3 and H4 (PubMed:18406326). Involved in endoderm differentiation via its association with long non-coding RNA (IncRNA) DIGIT: BRD3 undergoes liquid-liquid phase separation upon binding to IncRNA DIGIT, promoting binding to histone H3 acetylated at 'Lys-18' (H3K18ac) to induce endoderm gene expression (PubMed:32895492). Also binds non-histones acetylated proteins, such as GATA1 and GATA2: regulates transcription by promoting the binding of the transcription factor GATA1 to its targets (By similarity).

Cellular Location

Nucleus. Chromosome. Note=Detected on chromatin


Tissue Location Ubiquitous..

BRD3 Antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

BRD3 Antibody - middle region - Images



BRD3 Antibody - middle region (Al10087) in Human Liver tumor cells using Western Blot


Host: Rabbit

Target Name: BRD3

Sample Tissue: Liver tumor lysates

Antibody Dilution: 1.0µg/ml

BRD3 Antibody - middle region - Background







This is a rabbit polyclonal antibody against BRD3. It was validated on Western Blot by Abgent. At Abgent we manufacture rabbit polyclonal antibodies on a large scale (200-1000 products/month) of high throughput manner. Our antibodies are peptide based and protein family oriented. We usually provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).