

INSM2 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI10161**Specification****INSM2 antibody - middle region - Product Information**

Application	WB
Primary Accession	Q96T92
Other Accession	NM_032594 , NP_115983
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Rat, Pig, Guinea Pig, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 59kDa KDa

INSM2 antibody - middle region - Additional Information**Gene ID** 84684**Alias Symbol** IA-6, mlt1, IA6**Other Names**

Insulinoma-associated protein 2, Zinc finger protein IA-6, INSM2, IA6

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-INSM2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

INSM2 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

INSM2 antibody - middle region - Protein Information**Name** INSM2**Synonyms** IA6**Function**

May function as a growth suppressor or tumor suppressor in liver cells and in certain neurons.

Cellular Location

Cytoplasm. Nucleus.

Tissue Location

Expressed in heart, liver, skeletal muscle, kidney and pancreas, and, to a lesser extent, in brain, lung and spleen. In the pancreas, expressed in islet cells, including insulin- and glucagon-producing alpha- and beta-cells, but not in acinar cells (at protein level). Detected in adrenal glands, particularly in the deeper layer of the cortex (at protein level).

INSM2 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

