

Bcl2l2 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10180**Specification**

Bcl2l2 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P70345
Other Accession	NM_007537 , NP_031563
Reactivity	Human, Mouse, Rat, Pig, Horse, Bovine, Dog
Predicted Host	Rat, Pig, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 21kDa KDa

Bcl2l2 Antibody - C-terminal region - Additional Information**Gene ID** 12050**Alias Symbol** **AW048834, Bcl-w, Gtrgal2, Gtrosa41, bclw, c98****Other Names**

Bcl-2-like protein 2, Bcl2-L-2, Apoptosis regulator Bcl-W, c98, Bcl2l2, Bclw, Kiaa0271

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Bcl2l2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Bcl2l2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Bcl2l2 Antibody - C-terminal region - Protein Information**Name** Bcl2l2**Synonyms** Bclw, Kiaa0271**Function**

Promotes cell survival. Blocks dexamethasone-induced apoptosis. Mediates survival of postmitotic Sertoli cells by suppressing death-promoting activity of BAX.

Cellular Location

Mitochondrion membrane; Peripheral membrane protein. Note=Loosely associated with the

mitochondrial membrane in healthy cells. During apoptosis, tightly bound to the membrane (By similarity).


Tissue Location

Expressed in almost all myeloid cell lines and in a wide range of tissues, with highest levels in brain, colon, and salivary gland

Bcl2l2 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Bcl2l2 Antibody - C-terminal region - Images**WB Suggested Anti-Bcl2l2 Antibody**

Titration: 1. µg/ml

Positive Control: Mouse Testis