

STK17B Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10182**Specification**

STK17B Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q53QE7
Other Accession	NM_004226 , NP_004217
Reactivity	Human, Rat, Dog
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40kDa KDa

STK17B Antibody - C-terminal region - Additional Information

Alias Symbol **DRAK2**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 μ l of distilled water. Final Anti-STK17B antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions


STK17B Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

STK17B Antibody - C-terminal region - Protein Information**STK17B Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STK17B Antibody - C-terminal region - Images



Host:Rabbit

Target Name:STK17B

Sample Tissue: A549 Whole cell lysate

S

Antibody Dilution:3.µg/ml