

Ywhag antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10188**Specification**

Ywhag antibody - N-terminal region - Product Information

Application	WB
Primary Accession	P61982
Other Accession	NM_018871 , NP_061359
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Zebrafish, Chicken, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	28kDa KDa

Ywhag antibody - N-terminal region - Additional Information**Gene ID** 22628**Alias Symbol** 14-3-3gamma, D7Bwg1348e**Other Names**

14-3-3 protein gamma, 14-3-3 protein gamma, N-terminally processed, Ywhag

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Ywhag antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Ywhag antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Ywhag antibody - N-terminal region - Protein Information**Name** Ywhag**Function**

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner. Promotes inactivation of WDR24 component of the GATOR2 complex by binding to phosphorylated WDR24.

Cellular Location


Cytoplasm {ECO:0000250|UniProtKB:P68252}.

Ywhag antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ywhag antibody - N-terminal region - Images



WB Suggested Anti-Ywhag Antibody

Titration: 1. µg/ml

Positive Control: Mouse Brain