

ZNF365 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10262**Specification****ZNF365 antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	Q70YC5
Other Accession	NM_199451 , NP_955523
Reactivity	Human, Mouse, Rat, Horse, Bovine, Dog
Predicted	Pig, Bovine, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51kDa KDa

ZNF365 antibody - N-terminal region - Additional Information**Gene ID** 22891**Alias Symbol** UAN, Su48, ZNF365D**Other Names**

Protein ZNF365, Protein su48, ZNF365, KIAA0844

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-ZNF365 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF365 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF365 antibody - N-terminal region - Protein Information**Name** ZNF365**Synonyms** DBZ {ECO:0000303|PubMed:17389905}, KIAA0**Function**

Involved in the regulation of neurogenesis. Negatively regulates neurite outgrowth (PubMed:17389905). Involved in the morphogenesis of basket cells in the somatosensory cortex during embryogenesis. Involved in the positive regulation of oligodendrocyte differentiation during postnatal growth. Involved in dendritic arborization, morphogenesis of spine density dendrite, and establishment of postsynaptic dendrite density in cortical pyramidal neurons (By similarity). Involved in homologous recombination (HR) repair pathway. Required for proper resolution of DNA double-strand breaks

(DSBs) by HR. Is required for recovery of stalled replication forks, and directly contributes to genomic stability. Interacts with PARP1 and mediates MRE11-dependent DNA end resection during replication fork recovery (PubMed:23966166). Contributes to genomic stability by preventing telomere dysfunction (PubMed:23776040).

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=localizes to the centrosome at all stages of the cell cycle

Tissue Location

Isoform 1 is expressed in brain. Isoform 2 is expressed in placenta and at low level in lung and liver. Isoform 3 is expressed in kidney and pancreas. Isoform 1 is expressed exclusively in brain (PubMed:17389905).

ZNF365 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

