

Ywhae antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10309**Specification****Ywhae antibody - C-terminal region - Product Information**

Application	WB, IHC
Primary Accession	P62259
Other Accession	NM_009536 , NP_033562
Reactivity	Human, Mouse, Rat, Zebrafish, Sheep, Horse, Bovine
Predicted	Mouse, Zebrafish, Chicken, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29kDa KDa

Ywhae antibody - C-terminal region - Additional Information**Gene ID** 22627

Alias Symbol	AU019196
Other Names	
14-3-3 protein epsilon, 14-3-3E, Ywhae	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Ywhae antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Ywhae antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Ywhae antibody - C-terminal region - Protein Information**Name** Ywhae**Function**

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner. Positively regulates phosphorylated protein HSF1 nuclear export to the cytoplasm.

Cellular Location

Nucleus {ECO:0000250|UniProtKB:P62258}. Cytoplasm {ECO:0000250|UniProtKB:P62258}.

Melanosome {ECO:0000250|UniProtKB:P62258}

Ywhae antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)