

**KIN antibody - middle region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI10340****Specification****KIN antibody - middle region - Product Information**

Application	WB
Primary Accession	<a href="#">O60870</a>
Other Accession	<a href="#">NM_012311</a> , <a href="#">NP_036443</a>
Reactivity	Human, Mouse, Rat, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Chicken, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	45kDa KDa

**KIN antibody - middle region - Additional Information****Gene ID** 22944**Alias Symbol** BTCD, KIN17**Other Names**

DNA/RNA-binding protein KIN17, Binding to curved DNA, KIN, antigenic determinant of recA protein homolog, KIN ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=6327](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=6327) target="\_blank">HGNC:6327</a>), BTCD, KIN17

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 100 ul of distilled water. Final anti-KIN antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

KIN antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

**KIN antibody - middle region - Protein Information****Name** KIN ([HGNC:6327](#))**Synonyms** BTCD, KIN17**Function**

Involved in DNA replication and the cellular response to DNA damage. May participate in DNA replication factories and create a bridge between DNA replication and repair mediated by high molecular weight complexes. May play a role in illegitimate recombination and regulation of gene expression. May participate in mRNA processing. Binds, in vitro, to double-stranded DNA. Also

shown to bind preferentially to curved DNA in vitro and in vivo (By similarity). Binds via its C-terminal domain to RNA in vitro.

**Cellular Location**

Nucleus. Cytoplasm. Note=During S phase, strongly associated with the nuclear matrix, and to chromosomal DNA in the presence of DNA damage. Also shows cytoplasmic localization in elongated spermatids {ECO:0000250|UniProtKB:Q8K339, ECO:0000269|PubMed:11880372, ECO:0000269|PubMed:12359749, ECO:0000269|PubMed:12754299, ECO:0000269|PubMed:12853634, ECO:0000269|PubMed:15831485}

**Tissue Location**

Ubiquitously expressed in all tissues examined, with highest levels in skeletal muscle, heart and testis Differentially expressed in non-tumorigenic and tumorigenic cell lines Highly expressed in proliferating epithelial keratinocyte cells in vitro (at protein level).

**KIN antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)