

HEY1 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10371**Specification****HEY1 Antibody - C-terminal region - Product Information**

Application	WB, IHC
Primary Accession	O9Y5J3
Other Accession	NM_012258 , NP_036390
Reactivity	Human, Mouse, Rat, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Pig, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32kDa kDa

HEY1 Antibody - C-terminal region - Additional Information**Gene ID** 23462**Alias Symbol** BHLHb31, CHF2, HERP2, HESR1, HRT-1, OAF1**Other Names**

Hairy/enhancer-of-split related with YRPW motif protein 1, Cardiovascular helix-loop-helix factor 2, CHF-2, Class B basic helix-loop-helix protein 31, bHLHb31, HES-related repressor protein 1, Hairy and enhancer of split-related protein 1, HESR-1, Hairy-related transcription factor 1, HRT-1, hHRT1, HEY1, BHLHB31, CHF2, HERP2, HESR1, HRT1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-HEY1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

HEY1 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

HEY1 Antibody - C-terminal region - Protein Information**Name** HEY1**Synonyms** BHLHB31, CHF2, HERP2, HESR1, HRT1**Function**

Transcriptional repressor which binds preferentially to the canonical E box sequence 5'-CACGTG-3' (PubMed: <http://www.uniprot.org/citations/11095750>).

Downstream effector of Notch signaling required for cardiovascular development. Specifically required for the Notch-induced endocardial epithelial to mesenchymal transition, which is itself critical for cardiac valve and septum development. May be required in conjunction with HEY2 to specify arterial cell fate or identity. Promotes maintenance of neuronal precursor cells and glial versus neuronal fate specification. Represses transcription by the cardiac transcriptional activators GATA4 and GATA6 and by the neuronal bHLH factors ASCL1/MASH1 and NEUROD4/MATH3 (PubMed:15485867). Involved in the regulation of liver cancer cells self-renewal (PubMed:25985737).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00380, ECO:0000255|PROSITE-ProRule:PRU00981, ECO:0000269|PubMed:26068074}



Tissue Location

Expressed in the somitic mesoderm, the central nervous system, the kidney, the heart, nasal epithelium, and limbs

HEY1 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)




Host:Rabbit

Target Name:HEY1

Sample Tissue: Hela Whole cell lysate
s

Antibody Dilution:1.µg/ml



Rabbit Anti-HEY1 Antibody


Catalog Number: AI10371

Paraffin Embedded Tissue: Human Lung

Cellular Data: Alveolar cells

Antibody Concentration: 4.-8. µg/ml

Magnification: 4X



Rabbit Anti-HEY1 Antibody

Catalog Number: AI10371

Paraffin Embedded Tissue: Human Kidney

Cellular Data: Epithelial cells of renal tubule

Antibody Concentration: 4.-8. µg/ml

Magnification: 4X