

HMG20A antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI10415**Specification****HMG20A antibody - middle region - Product Information**

Application	WB
Primary Accession	Q9NP66
Other Accession	NM_018200 , NP_060670
Reactivity	Human, Mouse, Rat, Horse, Bovine, Dog
Predicted	Rat, Pig, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40kDa KDa

HMG20A antibody - middle region - Additional Information**Gene ID** 10363**Alias Symbol** **FLJ10739, HMGX1, HMGXB1****Other Names**

High mobility group protein 20A, HMG box-containing protein 20A, HMG domain-containing protein 1, HMG domain-containing protein HMGX1, HMG20A, HMGX1, HMGXB1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-HMG20A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

HMG20A antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

HMG20A antibody - middle region - Protein Information**Name** HMG20A**Synonyms** HMGX1, HMGXB1**Function**

Plays a role in neuronal differentiation as chromatin- associated protein. Acts as inhibitor of HMG20B. Overcomes the repressive effects of the neuronal silencer REST and induces the activation of neuronal-specific genes. Involved in the recruitment of the histone methyltransferase KMT2A/MLL1 and consequent increased methylation of histone H3 lysine 4 (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00267}.

Tissue Location

Ubiquitous..

HMG20A antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

