

PPP1R13L antibody - middle region

Rabbit Polyclonal Antibody Catalog # Al10420

Specification

PPP1R13L antibody - middle region - Product Information

Application WB, IHC Primary Accession Q8WUF5

Other Accession <u>NM 006663</u>, <u>NP 006654</u>

Reactivity Human, Mouse, Rat, Pig, Horse, Bovine,

Dog

Predicted Human, Mouse, Bovine, Guinea Pig, Dog

Host Rabbit
Clonality Polyclonal
Calculated MW 36kDa KDa

PPP1R13L antibody - middle region - Additional Information

Gene ID 10848

Alias Symbol

IASPP, NKIP1, RAI, iASPP gene, RAI4

Other Names

RelA-associated inhibitor, Inhibitor of ASPP protein, Protein iASPP, NFkB-interacting protein 1, PPP1R13B-like protein, PPP1R13L, IASPP, NKIP1, PPP1R13BL, RAI

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PPP1R13L antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

PPP1R13L antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

PPP1R13L antibody - middle region - Protein Information

Name PPP1R13L

Synonyms IASPP, NKIP1, PPP1R13BL, RAI

Function

Regulator that plays a central role in regulation of apoptosis and transcription via its interaction with NF-kappa-B and p53/TP53 proteins. Blocks transcription of HIV-1 virus by inhibiting the action of both NF-kappa-B and SP1. Also inhibits p53/TP53 function, possibly by preventing the association between p53/TP53 and ASPP1 or ASPP2, and therefore suppressing the subsequent activation of apoptosis (PubMed:<a href="http://www.uniprot.org/citations/12524540"





Tel: 858.875.1900 Fax: 858.875.1999

target=" blank">12524540). Is involved in NF-kappa-B dependent negative regulation of inflammatory response (PubMed:28069640).

Cellular Location

Cytoplasm. Nucleus Note=Predominantly cytoplasmic but also nuclear

Tissue Location

Highly expressed in heart, placenta and prostate. Weakly expressed in brain, liver, skeletal muscle, testis and peripheral blood leukocyte.

PPP1R13L antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture