

ARID3A antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI10430**Specification****ARID3A antibody - middle region - Product Information**

Application	WB
Primary Accession	Q99856
Other Accession	NM_005224 , NP_005215
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Horse, Bovine, Dog
Predicted Host	Rat, Pig, Bovine, Guinea Pig, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 63kDa KDa

ARID3A antibody - middle region - Additional Information**Gene ID 1820**

Alias Symbol **DRIL1, DRIL3, BRIGHT, E2FBP1**

Other Names

AT-rich interactive domain-containing protein 3A, ARID domain-containing protein 3A, B-cell regulator of IgH transcription, Bright, Dead ringer-like protein 1, E2F-binding protein 1, ARID3A, DRIL1, DRIL3, DRX, E2FBP1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ARID3A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ARID3A antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

ARID3A antibody - middle region - Protein Information

Name ARID3A

Synonyms DRIL1, DRIL3, DRX, E2FBP1

Function

Transcription factor which may be involved in the control of cell cycle progression by the RB1/E2F1 pathway and in B-cell differentiation.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00355, ECO:0000269|PubMed:17400556}.
Cytoplasm Note=Shuttles between nucleus and cytoplasm

Tissue Location

Widely expressed, with highest expression in skeletal muscle, thalamus, and colon

ARID3A antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)