

GATAD1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10476**Specification****GATAD1 antibody - C-terminal region - Product Information**

Application	WB
Primary Accession	Q8WUU5
Other Accession	NM_021167 , NP_066990
Reactivity	Human, Mouse, Rat, Zebrafish, Horse, Bovine, Dog
Predicted Host	Mouse, Rat, Chicken, Guinea Pig, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 29kDa KDa

GATAD1 antibody - C-terminal region - Additional Information**Gene ID** 57798**Alias Symbol** ODAG, RG083M05.2**Other Names**

GATA zinc finger domain-containing protein 1, Ocular development-associated gene protein, GATAD1, ODAG

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-GATAD1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

GATAD1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

GATAD1 antibody - C-terminal region - Protein Information**Name** GATAD1**Synonyms** ODAG**Function**

Component of some chromatin complex recruited to chromatin sites methylated 'Lys-4' of histone H3 (H3K4me), with a preference for trimethylated form (H3K4me3).

Cellular Location

Nucleus.

Tissue Location

Ubiquitously expressed among various tissue types. Expressed in left ventricular myocytes.

GATAD1 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

