

PHYHIP antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10574**Specification****PHYHIP antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	Q92561
Other Accession	NM_014759 , NP_055574
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Zebrafish, Pig, Chicken, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38kDa KDa

PHYHIP antibody - N-terminal region - Additional Information**Gene ID** 9796

Alias Symbol **DYRK1AP3, KIAA0273, PAHX-AP, PAHXAP1**

Other Names
Phytanoyl-CoA hydroxylase-interacting protein, Phytanoyl-CoA hydroxylase-associated protein 1, PAHX-AP1, PAHXAP1, PHYHIP, DYRK1AP3, KIAA0273

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PHYHIP antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

PHYHIP antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PHYHIP antibody - N-terminal region - Protein Information**Name** PHYHIP**Synonyms** DYRK1AP3, KIAA0273**Function**

Its interaction with PHYH suggests a role in the development of the central system.

Tissue Location

Highly expressed in the brain.

PHYHIP antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)