

KIF22 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10582**Specification****KIF22 antibody - C-terminal region - Product Information**

Application	WB
Primary Accession	Q14807
Other Accession	NM_007317 , NP_015556
Reactivity	Human, Mouse, Rat, Pig, Horse, Yeast, Bovine, Dog
Predicted	Human, Mouse, Rat, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	73kDa KDa

KIF22 antibody - C-terminal region - Additional Information**Gene ID** 3835**Alias Symbol** KID, KNSL4, OBP, OBP-1, OBP-2, SEMDJL2, A-328A3.2**Other Names**

Kinesin-like protein KIF22, Kinesin-like DNA-binding protein, Kinesin-like protein 4, KIF22, KID, KNSL4

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-KIF22 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

KIF22 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

KIF22 antibody - C-terminal region - Protein Information**Name** KIF22**Synonyms** KID, KNSL4**Function**

Kinesin family member that is involved in spindle formation and the movements of chromosomes during mitosis and meiosis. Binds to microtubules and to DNA (By similarity). Plays a role in congression of laterally attached chromosomes in NDC80-depleted cells (PubMed:25743205).

Cellular Location

Nucleus. Cytoplasm, cytoskeleton

Tissue Location

Expressed in bone, cartilage, joint capsule, ligament, skin, and primary cultured chondrocytes

KIF22 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)