

CXORF6 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10618**Specification****CXORF6 antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	Q13495
Other Accession	NM_005491 , NP_005482
Reactivity	Human, Mouse, Rat, Pig, Horse, Dog
Predicted	Human, Mouse, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	74kDa KDa

CXORF6 antibody - N-terminal region - Additional Information**Gene ID** 10046**Alias Symbol** CG1, F18, HYSP2, CXorf6**Other Names**

Mastermind-like domain-containing protein 1, F18, Protein CG1, MAMLD1, CG1, CXorf6

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-CXORF6 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CXORF6 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CXORF6 antibody - N-terminal region - Protein Information**Name** MAMLD1**Synonyms** CG1, CXorf6**Function**

Transactivates the HES3 promoter independently of NOTCH proteins. HES3 is a non-canonical NOTCH target gene which lacks binding sites for RBPJ.

Cellular Location

Nucleus. Note=Punctate nuclear localization

Tissue Location

Expressed in fetal brain, fetal ovary and fetal testis. Expressed in adult brain, ovary, skin, testis, uterus. Highly expressed in skeletal muscle.

CXORF6 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

