

6720456H20Rik Antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI10634**Specification**

6720456H20Rik Antibody - middle region - Product Information

Application	WB
Primary Accession	Q8BMD6
Other Accession	NM_172600 , NP_766188
Reactivity	Human, Mouse, Rat, Pig, Horse, Yeast, Bovine, Dog
Predicted Host	Human, Mouse, Rat, Pig, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 63kDa KDa

6720456H20Rik Antibody - middle region - Additional Information**Gene ID** 218989**Alias Symbol** 5930435P15, 6720456H20Rik
Other Names
Transmembrane protein 260, Tmem260**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-6720456H20Rik antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

6720456H20Rik Antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

6720456H20Rik Antibody - middle region - Protein Information**Name** Tmem260 {ECO:0000312|MGI:MGI:2443219}**Function**

O-mannosyl-transferase that transfers mannosyl residues to the hydroxyl group of serine or threonine residues of proteins. Specifically glycosylates the IPT/TIG domain of target proteins, such as MET and MST1R/RON. TMEM260-mediated O-mannosylated residues are composed of single mannose glycans that are not elongated or modified.

Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q9NX78}; Multi-pass membrane


protein

6720456H20Rik Antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

6720456H20Rik Antibody - middle region - Images



Host:Rabbit

Target Name:672456H2Rik

Sample Tissue: Mouse Testis lysates

Antibody Dilution:1.µg/ml