

COLQ antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10635**Specification**

COLQ antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q6DK18
Other Accession	NM_005677 , NP_005668
Reactivity	Human, Mouse, Rat, Horse, Bovine
Predicted	Human, Mouse, Rat, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48kDa KDa

COLQ antibody - N-terminal region - Additional Information

Alias Symbol **EAD**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-COLQ antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


COLQ antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

COLQ antibody - N-terminal region - Protein Information**COLQ antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

COLQ antibody - N-terminal region - Images

**WB Suggested Anti-COLQ Antibody****Titration: 1.25 µg/ml**

Positive Control: HepG2 Whole Cell

COLQ antibody - N-terminal region - References

Ting, A.K., et al., (2005) Chem. Biol. Interact. 157-158, 63-70
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.