

HCFC2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10654**Specification**

HCFC2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9Y5Z7
Other Accession	NM_013320 , NP_037452
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	87kDa KDa

HCFC2 antibody - C-terminal region - Additional Information**Gene ID** 29915

Alias Symbol	HCF-2, HCF2
Other Names	
Host cell factor 2, HCF-2, C2 factor, HCFC2	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-HCFC2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

HCFC2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

HCFC2 antibody - C-terminal region - Protein Information**Name** HCFC2**Cellular Location**

Cytoplasm. Nucleus

Tissue Location


Highly expressed in testis. Detected at lower levels in spleen, thymus, prostate, ovary, small intestine and colon

HCFC2 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HCFC2 antibody - C-terminal region - Images



WB Suggested Anti-HCFC2 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Jurkat cell lysate

HCFC2 antibody - C-terminal region - References

Yokoyama, A., (2004) Mol. Cell. Biol. 24 (13), 5639-5649 Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.