

DKFZP761C169 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10655**Specification**

DKFZP761C169 antibody - N-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q86WP2
Other Accession	NM_022913 , NP_075064
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Rat, Chicken, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 52kDa KDa

DKFZP761C169 antibody - N-terminal region - Additional Information**Gene ID** 65056**Alias Symbol** GPBP, SSH6, VASCULIN, DKFZP761C169
Other Names
Vasculin, GC-rich promoter-binding protein 1, Vascular wall-linked protein, GPBP1, GPBP, SSH6**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-DKFZP761C169 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

DKFZP761C169 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DKFZP761C169 antibody - N-terminal region - Protein Information**Name** GPBP1**Synonyms** GPBP, SSH6**Function**

Functions as a GC-rich promoter-specific transactivating transcription factor.

Cellular Location

Nucleus. Cytoplasm. Note=According to PubMed:12842993, it localizes to the cytoplasm.


Tissue Location

Widely expressed. Some isoforms may be specifically expressed in veins and arteries (at protein level). Isoform 4 is widely expressed. Isoform 1, isoform 2 and isoform 3 may be specifically expressed in vascular smooth muscle cells


DKFZP761C169 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DKFZP761C169 antibody - N-terminal region - Images

Human kidney



WB Suggested Anti-DKFZP761C169 Antibody Titration: 0.2-1 µg/ml
Positive Control: HepG2 cell lysate

DKFZP761C169 antibody - N-terminal region - References

Hsu, L.C., et al., (2003) Kidd, V.J., Mol. Cell. Biol. 23 (23), 8773-8785
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.