

CORO1A antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10667**Specification**

CORO1A antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q2YD73
Other Accession	NM_007074 , NP_009005
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Rabbit, Pig, Horse, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 51kDa KDa

CORO1A antibody - C-terminal region - Additional Information

Alias Symbol **p57, TACO, CLABP, HCORO1, CLIPINA**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-CORO1A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


CORO1A antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CORO1A antibody - C-terminal region - Protein Information**CORO1A antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CORO1A antibody - C-terminal region - Images



WB Suggested Anti-CORO1A Antibody Titration: 1.25µg/ml

ELISA Titer: 1:1562500

Positive Control: Jurkat cell lysate

CORO1A is supported by BioGPS gene expression data to be expressed in Jurkat

CORO1A antibody - C-terminal region - References

Anand,P.K., et al., (2005) FEMS Microbiol. Lett. 250 (1), 137-144
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.