

BTBD14A antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10700**Specification**

BTBD14A antibody - N-terminal region - Product Information

Application	WB, IHC
Primary Accession	Q96BF6
Other Accession	NM_144653 , NP_653254
Reactivity	Human, Mouse, Rat, Horse, Bovine, Dog
Predicted	Human, Chicken, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	63kDa KDa

BTBD14A antibody - N-terminal region - Additional Information**Gene ID** 138151**Alias Symbol** **BTBD14, MGC23427, BEND9, BTBD14A**
Other Names

Nucleus accumbens-associated protein 2, NAC-2, BTB/POZ domain-containing protein 14A, Repressor with BTB domain and BEN domain, NACC2, BTBD14A, NAC2, RBB

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-BTBD14A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

BTBD14A antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

BTBD14A antibody - N-terminal region - Protein Information**Name** NACC2**Synonyms** BTBD14A, NAC2, RBB**Function**

Functions as a transcriptional repressor through its association with the NuRD complex. Recruits the NuRD complex to the promoter of MDM2, leading to the repression of MDM2 transcription and subsequent stability of p53/TP53.


Cellular Location

Nucleus. Note=Predominantly associated with chromatin


BTBD14A antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BTBD14A antibody - N-terminal region - Images

Human kidney



WB Suggested Anti-BTBD14A Antibody Titration: 0.625µg/ml
Positive Control: Jurkat cell lysate