

Hpcal1 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10706**Specification**

Hpcal1 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P62748
Other Accession	NM_016677 , NP_057886
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Sheep, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Zebrafish, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22kDa KDa

Hpcal1 Antibody - C-terminal region - Additional Information**Gene ID** 53602**Alias Symbol** NVP-3, Nvp3, VILIP3, VnsI3, VsnI3**Other Names**

Hippocalcin-like protein 1, Neural visinin-like protein 3, NVL-3, NVP-3, Visinin-like protein 3, VILIP-3, Hpcal1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Hpcal1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Hpcal1 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Hpcal1 Antibody - C-terminal region - Protein Information**Name** Hpcal1**Function**

May be involved in the calcium-dependent regulation of rhodopsin phosphorylation.

Cellular Location


Membrane {ECO:0000250|UniProtKB:P37235}; Lipid- anchor {ECO:0000250|UniProtKB:P37235}

Hpcal1 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Hpcal1 Antibody - C-terminal region - Images



WB Suggested Anti-Hpcal1 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Stomach