

ASPH antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI10707**Specification**

ASPH antibody - middle region - Product Information

Application	IHC, WB
Primary Accession	O9H291
Other Accession	NM_032468 , NP_115857
Reactivity	Human, Mouse, Rat, Pig, Bovine
Predicted	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34kDa KDa

ASPH antibody - middle region - Additional Information

Alias Symbol **BAH, CASQ2BP1, HAAH, JCTN, junctin, AAH**
Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ASPH antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


ASPH antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

ASPH antibody - middle region - Protein Information**ASPH antibody - middle region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ASPH antibody - middle region - Images



Immunohistochemistry with Small intestine, myenteric plexus tissue at an antibody concentration of 5 µg/ml using anti-ASPH antibody (AI10707)



Rabbit Anti-ASPH Antibody

Catalog Number: AI10707

Formalin Fixed Paraffin Embedded Tissue: Human Adult heart Observed Staining: Cytoplasmic, Membrane

Primary Antibody

Concentration: 1:600

Secondary Antibody: Donkey anti-Rabbit-Cy2/3


Secondary Antibody

Concentration: 1:200

Magnification: 20X

Exposure Time: 0.5 – 2.0 sec

Protocol located in Reviews and Data.



WB Suggested Anti-ASPH Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:62500

Positive Control: ACHN cell lysate

ASPH antibody - middle region - References

Gerhard,D.S., (2008) Oncol. Rep. 19 (3), 795-800 Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.