

Thra antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10730**Specification**

Thra antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P63059
Other Accession	NM_001017960 , NP_001017960
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Sheep, Horse, Bovine, Dog
Predicted Host	Mouse, Rat, Rabbit, Zebrafish, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 55kDa KDa

Thra antibody - C-terminal region - Additional Information**Gene ID** 81812**Alias Symbol** **ERBA1, Thra1****Other Names**

Thyroid hormone receptor alpha, Nuclear receptor subfamily 1 group A member 1, c-erbA-1, c-erbA-alpha, Thra, C-erba-alpha, Nr1a1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Thra antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Thra antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Thra antibody - C-terminal region - Protein Information**Name** Thra**Synonyms** C-erba-alpha, Nr1a1**Function**

Nuclear hormone receptor that can act as a repressor or activator of transcription. High affinity receptor for thyroid hormones, including triiodothyronine and thyroxine.

Cellular Location


Nucleus.

Thra antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Thra antibody - C-terminal region - Images



WB Suggested Anti-Thra Antibody Titration: 1.0 µg/ml
Positive Control: Rat Muscle