

ZNF396 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI10746**Specification**

ZNF396 antibody - middle region - Product Information

Application	WB
Primary Accession	O96N95
Other Accession	NM_145756 , NP_665699
Reactivity	Human, Rabbit
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38kDa KDa

ZNF396 antibody - middle region - Additional Information**Gene ID** 252884**Alias Symbol** **FLJ31213, ZSCAN14****Other Names**

Zinc finger protein 396, Zinc finger and SCAN domain-containing protein 14, ZNF396, ZSCAN14

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZNF396 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF396 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF396 antibody - middle region - Protein Information**Name** ZNF396**Synonyms** ZSCAN14**Function**

Isoform 1 and isoform 2 act as DNA-dependent transcriptional repressors.

Cellular Location

[Isoform 1]: Nucleus.

Tissue Location

Expressed strongly in liver, moderately in skeletal muscle and weakly in kidney, pancreas, spleen


and prostate

ZNF396 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF396 antibody - middle region - Images



WB Suggested Anti-ZNF396 Antibody Titration: 0.2-1 $\mu\text{g/ml}$
ELISA Titer: 1:62500
Positive Control: Jurkat cell lysate

ZNF396 antibody - middle region - References

Wu, Y., Gene 310, 193-201 (2003) Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.