

CLIC5 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10811**Specification**

CLIC5 antibody - C-terminal region - Product Information

Application	WB, IHC
Primary Accession	Q53G01
Other Accession	NM_016929 , NP_058625
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Rat, Pig, Chicken
Clonality	Rabbit
Calculated MW	Polyclonal 28kDa KDa

CLIC5 antibody - C-terminal region - Additional Information

Alias Symbol **MST130, MSTP130**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CLIC5 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


CLIC5 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CLIC5 antibody - C-terminal region - Protein Information**CLIC5 antibody - C-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CLIC5 antibody - C-terminal region - Images



Human Heart



WB Suggested Anti-CLIC5 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:1562500

Positive Control: Jurkat cell lysate

There is BioGPS gene expression data showing that CLIC5 is expressed in Jurkat

CLIC5 antibody - C-terminal region - References

Berryman, M., et al., (2004) J. Biol. Chem. 279 (33), 34794-34801
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.