

KCNK10 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10819**Specification**

KCNK10 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P57789
Other Accession	NM_021161 , NP_066984
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rabbit, Pig, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59kDa kDa

KCNK10 antibody - C-terminal region - Additional Information**Gene ID** 54207**Alias Symbol** TREK2, TREK-2, K2p10.1**Other Names**

Potassium channel subfamily K member 10, Outward rectifying potassium channel protein TREK-2, TREK-2 K(+) channel subunit, KCNK10, TREK2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-KCNK10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

KCNK10 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

KCNK10 antibody - C-terminal region - Protein Information**Name** KCNK10**Synonyms** TREK2**Function**

Outward rectifying potassium channel. Produces rapidly activating and non-inactivating outward rectifier K(+) currents. Activated by arachidonic acid and other naturally occurring unsaturated free fatty acids.

Cellular Location

Membrane; Multi-pass membrane protein


Tissue Location

Abundantly expressed in pancreas and kidney and to a lower level in brain, testis, colon, and small intestine. Isoform b is strongly expressed in kidney (primarily in the proximal tubule) and pancreas, whereas isoform c is abundantly expressed in brain

KCNK10 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KCNK10 antibody - C-terminal region - Images

WB Suggested Anti-KCNK10 Antibody Titration: 2.5µg/ml

ELISA Titer: 1:62500

Positive Control: Jurkat cell lysate

KCNK10 antibody - C-terminal region - References

Gu, W., et al., (2002) J. Physiol. (Lond.) 539 (PT 3), 657-668
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.