

VGLL1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10907**Specification**

VGLL1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q99990
Other Accession	NM_016267 , NP_057351
Reactivity	Human, Rat
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29kDa KDa

VGLL1 antibody - N-terminal region - Additional Information**Gene ID** 51442

Alias Symbol	TDU, VGL1
Other Names	
Transcription cofactor vestigial-like protein 1, Vgl-1, Protein TONDU, VGLL1, TDU	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-VGLL1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

VGLL1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

VGLL1 antibody - N-terminal region - Protein Information**Name** VGLL1**Synonyms** TDU**Function**

May act as a specific coactivator for the mammalian TEFs.

Cellular Location


Nucleus.

VGLL1 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

VGLL1 antibody - N-terminal region - Images



WB Suggested Anti-VGLL1 Antibody Titration: 2.5µg/ml

ELISA Titer: 1:312500

Positive Control: Jurkat cell lysate

VGLL1 antibody - N-terminal region - References

Vaudin, P., et al., (1999) Development 126 (21), 4807-4816
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.