

ZNF677 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10964**Specification**

ZNF677 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q86XU0
Other Accession	NM_182609 , NP_872415
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	68kDa KDa

ZNF677 antibody - C-terminal region - Additional Information**Gene ID** 342926

Alias Symbol	MGC48625
Other Names	
Zinc finger protein 677, ZNF677	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZNF677 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF677 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF677 antibody - C-terminal region - Protein Information**Name** ZNF677**Function**

May be involved in transcriptional regulation.

Cellular Location


Nucleus.

ZNF677 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF677 antibody - C-terminal region - Images



WB Suggested Anti-ZNF677 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: Human Placenta

ZNF677 antibody - C-terminal region - References

Gerhard,D.S., Genome Res. 14 (10B), 2121-2127 (2004)Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.