

CX40.1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11008**Specification**

CX40.1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8N2R7
Other Accession	NM_153368 , NP_699199
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41kDa KDa

CX40.1 antibody - C-terminal region - Additional Information

Alias Symbol **CX40.1, RP11-425A6.2**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-CX40.1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


CX40.1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CX40.1 antibody - C-terminal region - Protein Information**CX40.1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CX40.1 antibody - C-terminal region - Images



90 kDa
60 kDa
42 kDa
32 kDa
23 kDa

WB Suggested Anti-CX40.1 Antibody Titration: 0.6µg/ml

ELISA Titer: 1:62500

Positive Control: HepG2 cell lysate

CX40.1 antibody - C-terminal region - References

S, et al., (2003) Cell Commun. Adhes. 10 (1), 27-36
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.