

Foxd2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11069**Specification**

Foxd2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O35392
Other Accession	NM_008593 , NP_032619
Reactivity	Human, Mouse, Rat, Horse, Bovine
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	49 KDa

Foxd2 antibody - C-terminal region - Additional Information**Gene ID** 17301**Alias Symbol** **Mf2, AI426778****Other Names**

Forkhead box protein D2, Mesoderm/mesenchyme forkhead 2, MF-2, Foxd2, Mf2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Foxd2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Foxd2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Foxd2 antibody - C-terminal region - Protein Information**Name** Foxd2**Synonyms** Mf2**Function**

Probable transcription factor involved in embryogenesis and somatogenesis.

Cellular Location

Nucleus.

Tissue Location

Expressed at high levels in the ventral region of newly formed somites, in sclerotomal derivatives,


in lateral plate and cephalic mesoderm and in the first and second branchial arches. Other regions of mesodermal expression include the developing tongue, meninges, nose, whiskers, kidney, genital tubercle and limb joints. In the nervous system it is transcribed in restricted regions of the mid- and forebrain.

Foxd2 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Foxd2 antibody - C-terminal region - Images



WB Suggested Anti-Foxd2 Antibody Titration: 0.2-1 µg/ml
Positive Control: Mouse Thymus