

Nab2 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI11072**Specification**

Nab2 antibody - middle region - Product Information

Application	WB
Primary Accession	Q61127
Other Accession	NM_008668 , NP_032694
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57 KDa

Nab2 antibody - middle region - Additional Information**Gene ID** 17937**Alias Symbol** **AI451907****Other Names**

NGFI-A-binding protein 2, EGR-1-binding protein 2, Nab2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Nab2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Nab2 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Nab2 antibody - middle region - Protein Information**Name** Nab2**Function**

Acts as a transcriptional repressor for zinc finger transcription factors EGR1 and EGR2. Isoform 2 lacks repression ability.

Cellular Location

Nucleus. Note=Isoform 2 is not localized to the nucleus

Tissue Location


Highly expressed in brain and thymus, and at lower levels in spleen, kidney, heart and testis.
Isoform 1 is predominantly expressed in testis, whereas isoform 3 is more abundant in thymus

Nab2 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Nab2 antibody - middle region - Images



WB Suggested Anti-Nab2 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Mouse Kidney