

**Prrx2 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI11081****Specification**

---

**Prrx2 antibody - N-terminal region - Product Information**

|                   |                                                       |
|-------------------|-------------------------------------------------------|
| Application       | WB                                                    |
| Primary Accession | <a href="#">Q06348</a>                                |
| Other Accession   | <a href="#">NM_009116</a> , <a href="#">NP_033142</a> |
| Reactivity        | Mouse, Rat                                            |
| Predicted         | Mouse, Rat                                            |
| Host              | Rabbit                                                |
| Clonality         | Polyclonal                                            |
| Calculated MW     | 26kDa KDa                                             |

**Prrx2 antibody - N-terminal region - Additional Information****Gene ID** 20204**Alias Symbol** Prx2, S8**Other Names**

Paired mesoderm homeobox protein 2, PRX-2, Homeobox protein S8, Prrx2, Prx2, S8

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-Prrx2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

Prrx2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**Prrx2 antibody - N-terminal region - Protein Information****Name** Prrx2**Synonyms** Prx2, S8**Cellular Location**

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000255|PROSITE-ProRule:PRU00138}

**Tissue Location**


In the embryo, expressed in craniofacial mesenchyme, limb, heart, somites and sclerotomes. Absent from central and peripheral nervous systems, splanchnopleure and ectodermal derivatives

## Prrx2 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Prrx2 antibody - N-terminal region - Images



WB Suggested Anti-Prrx2 Antibody Titration: 0.2-1  $\mu$ g/ml  
Positive Control: Mouse Spleen