

# Ahrr antibody - middle region

Rabbit Polyclonal Antibody Catalog # Al11098

## **Specification**

# Ahrr antibody - middle region - Product Information

Application WB
Primary Accession Q3U1U7

Other Accession NM\_009644, NP\_033774

Reactivity
Predicted
Host
Clonality
Calculated MW

Mouse, Rat
Rabbit
Polyclonal
78kDa KDa

### Ahrr antibody - middle region - Additional Information

**Gene ID 11624** 

Alias Symbol KIAA1234, mKIAA1234

**Other Names** 

Aryl hydrocarbon receptor repressor, AhR repressor, AhRR, Ahrr, Kiaa1234

#### **Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

### **Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-Ahrr antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

### **Precautions**

Ahrr antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

# Ahrr antibody - middle region - Protein Information

Name Ahrr

Synonyms Kiaa1234

#### **Function**

Mediates dioxin toxicity and is involved in regulation of cell growth and differentiation. Represses the transcription activity of AHR by competing with this transcription factor for heterodimer formation with the ARNT and subsequently binding to the xenobiotic response element (XRE) sequence present in the promoter regulatory region of variety of genes. Represses CYP1A1 by binding the XRE sequence and recruiting ANKRA2, HDAC4 and/or HDAC5. Autoregulates its expression by associating with its own XRE site.



## **Cellular Location**


Cytoplasm. Nucleus {ECO:0000255|PROSITE-ProRule:PRU00981, ECO:0000269|PubMed:9887096} Note=Predominantly in the nuclear compartment. First cytoplasmic, translocates into the nuclear compartment upon interaction with ARNT in the cytoplasmic compartment

# **Ahrr antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Ahrr antibody - middle region - Images



WB Suggested Anti-Ahrr Antibody Titration: 0.2-1 µg/ml

Positive Control: Mouse Muscle