

Msc antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11125**Specification**

Msc antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O88940
Other Accession	NM_010827 , NP_034957
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Dog
Predicted Host	Rat, Rabbit, Chicken, Horse, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 21kDa KDa

Msc antibody - C-terminal region - Additional Information**Gene ID** 17681

Alias Symbol	MyoR, bHLHa22
Other Names	
Musculin, Myogenic repressor, Msc, Myor	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Msc antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Msc antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Msc antibody - C-terminal region - Protein Information**Name** Msc**Synonyms** Myor**Function**

Transcription repressor that blocks myogenesis and activation of E-box dependent muscle genes.

Cellular Location


Nucleus.

Msc antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Msc antibody - C-terminal region - Images



WB Suggested Anti-Msc Antibody Titration: 0.2-1 µg/ml
Positive Control: Mouse Small Intestine