

NFATC2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11130**Specification**

NFATC2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q60591-3
Other Accession	NM_010899 , NP_035029
Reactivity	Human, Mouse, Rat, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	102kDa KDa

NFATC2 antibody - N-terminal region - Additional Information

Alias Symbol **NFAT1, Nfatp, NF-ATc2, NFAT1-D, AI607462**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-NFATC2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


NFATC2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NFATC2 antibody - N-terminal region - Protein Information**NFATC2 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NFATC2 antibody - N-terminal region - Images



WB Suggested Anti-NFATC2 Antibody Titration: 1.25µg/ml

ELISA Titer: 1:312500

Positive Control: SP2/0 cell lysate

NFATC2 antibody - N-terminal region - References

Glud, S.Z., et al., (2005) Blood 106 (10), 3546-3552 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles. Publications: Heer, R. et al. Phenotypic modulation of human urinary tract stroma-derived fibroblasts by transforming growth factor beta3. Urology 76, 509.e13-20 (2010). WB, IP, Mouse, Human, Rat, H, Bovine, Dog, Pig 20546875 Nagamoto-Combs, K. & Combs, C. K. Microglial phenotype is regulated by activity of the transcription factor, NFAT (nuclear factor of activated T cells). J. Neurosci. 30, 9641-6 (2010). WB, IP, Mouse, Human, Rat, H, Bovine, Dog, Pig 20631193