

Zscan12 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11155**Specification**

Zscan12 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9Z1D7
Other Accession	NM_016684 , NP_057893
Reactivity	Mouse, Rat
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57kDa KDa

Zscan12 antibody - C-terminal region - Additional Information**Gene ID** 22758**Alias Symbol** 2510038J07Rik, FPM315, Zfp96, mKIAA0426**Other Names**

Zinc finger and SCAN domain-containing protein 12, Zinc finger protein 96, Zfp-96, Zscan12, Zfp96, Znf96

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Zscan12 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Zscan12 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Zscan12 antibody - C-terminal region - Protein Information**Name** Zscan12**Synonyms** Zfp96, Znf96**Function**

May be involved in transcriptional regulation.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00187}.


Tissue Location

Testis specific..

Zscan12 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Zscan12 antibody - C-terminal region - Images

WB Suggested Anti-Zscan12 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Mouse Thymus