

IRX6 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11182**Specification**

IRX6 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8BFT1
Other Accession	NM_022428 , NP_071873
Reactivity	Mouse, Rat
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48kDa KDa

IRX6 antibody - C-terminal region - Additional Information**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-IRX6 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


IRX6 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

IRX6 antibody - C-terminal region - Protein Information**IRX6 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IRX6 antibody - C-terminal region - Images



WB Suggested Anti-IRX6 Antibody Titration: 1.25 μ g/ml

ELISA Titer: 1:12500

Positive Control: NIH/3T3 cell lysate

IRX6 antibody - C-terminal region - References

Pan, L., et al., (2005) Development 132 (4), 703-712
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.