

DMRTC2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11202**Specification**

DMRTC2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8°CGW9
Other Accession	NM_027732 , NP_082008
Reactivity	Mouse, Rat
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41kDa KDa

DMRTC2 antibody - N-terminal region - Additional Information

Alias Symbol **Dmrt7, 4933432E21Rik**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-DMRTC2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


DMRTC2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DMRTC2 antibody - N-terminal region - Protein Information**DMRTC2 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DMRTC2 antibody - N-terminal region - Images



WB Suggested Anti-DMRTC2 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: Human Testis

DMRTC2 antibody - N-terminal region - References

Kim, S., et al., (2003) Gene Expr. Patterns 3 (1), 77-82
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.